2

In the Claims:

Claim 1. (Currently Amended) A method for feeding an infant with milk equivalent infant formula, the method comprising:

- (a) feeding the infant by fore-milk equivalent infant formula; and
- (b) feeding the infant by hind-milk equivalent <u>infant formula</u>, wherein said feeding by fore-milk equivalent infant formula and said feeding by hind-milk infant formula equivalent are done sequentially during the same feeding session.

Claim 2. (Currently Amended) The method of claim 1, wherein said fore-milk equivalent infant formula has a volume of 30-60% of a total meal fed to the infant and a percentage of fat of 2.5 - 3.5%, and said hind-milk equivalent infant formula has a volume of 40 - 70% of the total meal and a percentage of fat of 3.7 - 5.5%.

Claim 3. (Currently Amended) The method of claim 1, wherein said feeding is performed either by feeding said fore-milk equivalent infant formula and said hind-milk equivalent infant formula one after another or by first feeding said fore-milk equivalent infant formula which is gradually admixed with said hind-milk equivalent infant formula.

Claim 4. (Currently Amended) The method of claim 2, wherein the percentage of said fore-milk equivalent infant formula in the total meal volume is 31 - 47% and the fat content therein is 2.8% - 3.3%, and the percentage of said hind-milk equivalent infant formula in the total meal volume is 53 - 67% and the fat content therein is 4.5% - 5%.

Claim 5. (Currently Amended) The method of claim 4, wherein the percentage of said fore-milk equivalent <u>infant formula</u> in the total meal is 40 % and the fat content therein is 3 %, and the percentage of said hind-milk equivalent <u>infant formula</u> in the total meal volume is 60 % and the fat content therein is 4.8 %.

Claim 6. (Currently Amended) The method of claim 1, wherein said feeding is performed using two bottles, one being filled with said fore-milk equivalent

3

infant formula and one being filled with said hind-milk equivalent infant formula, the contents of said bottles being forwarded one after the other.

Claim 7. (Currently Amended) The method of claim 1, wherein said feeding is performed with a bottle which is divided into two compartments, one being filled with said fore-milk equivalent <u>infant formula</u> and the other being filled with said hind-milk equivalent <u>infant formula</u>, the bottle being designed in such a manner enabling the infant to consume at first said fore-milk equivalent <u>infant formula</u> and thereafter said fore-milk equivalent <u>infant formula</u> gradually admixed with said hind-milk equivalent <u>infant formula</u>.

Claim 8. (Withdrawn) A kit for feeding an infant with milk equivalent, the kit comprising:

- (a) a first container containing a fore-milk equivalent formula; and
- (b) a second container containing a hind-milk equivalent formula.

Claim 9. (Withdrawn) The kit of claim 8, wherein said fore-milk equivalent formula has a percentage of fat of 2.5 % - 3.5 %, and said hind-milk equivalent formula has a percentage of fat of 3.7 % - 5.5 %.

Claim 10. (Withdrawn) The kit of claim 8, wherein said fore-milk equivalent formula has a percentage of fat of 2.8 - 3.3 %, and said hind-milk equivalent formula has a percentage of fat of 4.5 - 5 %.

Claim 11. (Withdrawn) The kit of claim 8, wherein said fore-milk equivalent formula has a percentage of fat of 3 %, and said hind-milk equivalent formula has a percentage of fat of 4.8 %.

Claim 12. (Withdrawn) The kit of claim 8, wherein an amount of said hind-milk equivalent formula in the kit is between 1.5 and 2.3 fold an amount of said fore-milk equivalent formula in the kit.

4

Claim 13. (Withdrawn) The kit of claim 8, wherein said foremilk equivalent formula and said hind-milk equivalent formula is each independently selected from the group consisting of a soy base milk equivalent formula and milk base milk equivalent formula.

Claim 14. (Withdrawn) The kit of claim 8, wherein said fore-milk equivalent formula and said hind-milk equivalent formula is each independently selected from the group consisting of a powder milk equivalent and a liquid milk equivalent formula.

Claim 15. (New) The method of claim 1, wherein said feeding the infant by fore-milk equivalent is done prior to said feeding the infant by hind-milk equivalent.